1

Neocallichirus rodfeldmanni, a proposed replacement name for Neocallichirus manningi Schweitzer, Feldmann, Fam, Hessin, Hetrick, Nyborg & Ross, 2003 (Crustacea: Decapoda: Axiidea: Callianassidae)

Matúš Hyžný

Department of Geology and Palaeontology, Faculty of Natural Sciences, Comenius University, Mlynská Dolina G1, Bratislava 842 15, Slovakia<hyzny.matus@gmail.com>

Key words: Decapoda, Axiidea, Callianassidae, Neocallichirus, replacement name

The fossil species *Neocallichirus manningi* Schweitzer, Feldmann, Fam, Hessin, Hetrick, Nyborg and Ross, 2003 was described from the Campanian strata of Canada (British Columbia). However, this name is already preoccupied with the extant species *Neocallichirus manningi* Kazmi and Kazmi, 1992, known from the Indian Ocean (Arabian Sea, Pakistan). Therefore a new replacement name *Neocallichirus rodfeldmanni* is proposed here for the fossil species *Neocallichirus manningi* Schweitzer, Feldmann, Fam, Hessin, Hetrick, Nyborg and Ross, 2003. The new replacement name honours Rodney M. Feldmann (Kent State University, Ohio, USA) who has contributed much to our knowledge of fossil decapod crustaceans.

Acknowledgement

I thank C. E. Schweitzer (Kent State University, Ohio, USA) for her encouragement.

References

Kazmi, Q. B. and Kazmi, M. A. (1992), A new species of a callianassid shrimp, *Neocallichirus manningi*, with a note on the genus *Neocallichirus* Sakai, 1988, not previously recorded from the Arabian Sea. *Crustaceana*, 63, 296–300.

Schweitzer, C. E., Feldmann, R. M., Fam, J., Hessin, W. A., Hetrick, S. W., Nyborg, T. G. and Ross, R. L. M. (2003), Cretaceous and Eocene decapod crustaceans from Southern Vancouver Island, British Columbia, Canada. NRC Research Press, Ottawa, Ontario, 66 p.

Accepted on September 11, 2009